

# Assessment of Nutritional Potency and Pomological Traits in the Wild and Cultivated Varieties of *Momordica charantia* L. (Cucurbitaceae)

Salma Suhara Beevy and Noor Muhammed Haseena Bai

Department of Botany, University of Kerala Kariavattom, Thiruvananthapuram  
695 508, India

## ABSTRACT

Nutrient composition in the fruits of cultivated and wild varieties of *M. charantia* viz, *M.charantia* var. *charantia* and var. *muricata* were evaluated to prioritize the wild accessions suitable for crop improvement programme. The nutritive value of the varieties were assessed in terms of the quantity of proteins, carbohydrate, vitamins, total sugar, reducing and non-reducing sugar and minerals like Na, Ca, Mg, Mn, Fe, Cu and Zn. The wild variety was found superior than the cultivar in terms of their nutrient content. The study recognized MC12, as the elite germplasm with highest concentration of Vit. E, B2, Cu and Ca and the accession MC10 as a rich source of carbohydrate, non-reducing sugar and Vit. B1. The accessions MC2, MC7, and MC15 were superior having higher concentration of protein, Vit. C and reducing sugar respectively. Correlation analysis was carried out to determine the relationship between fruit traits and nutrient composition. UPGMA dendrogram analysis suggests that the nutritional variation in the accessions was not determined by the fruit and seed morphology. However, a direct relationship between the nutrient contents and the fruit traits viz, fruit weight, length and fruit diameter was observed. The study recommends the accessions like MC12, MC7, MC10, and MC15 with high nutrient composition for bitter gourd breeding programme. The study emphasizes the conservation of the wild species, the rich sources of nutrients for crop improvement programme.

*Key words:* fruit traits, intra specific, *M. charantia* var.*charantia*, *M.charantia* var.*muricata*, nutrient contents.

## INTRODUCTION

*Momordica charantia* L. commonly known as bitter gourd, or balsam pear or bitter melon belonging to the family Cucurbitaceae is the most widely cultivated species of the genus *Momordica*. The species has many uses; the immature fruit, young stems, leaves and flowers are taken as vegetable, primarily in Asia and particularly in India (Reyce *et al.* 1994). *M. charantia* includes both the wild and cultivated varieties, and are distinguished by their nature of leaves and fruits (Decker-Walters, 1999; Behara *et al.*, 2006). The species possess two varieties viz: variety *charantia* which produces large fusiform fruits and variety *muricata* (wild) with small and round fruits (Chakravathy, 1990). The wild species offers great resources for breeding of the cultivated bitter gourd for the desirable edible or qualitative traits such as non bitterness, tolerance to abiotic stresses and resistance to several insect pests. According to a FAO report, at least one billion people are thought to use wild plants in their diet and for medicines. In most of the report it was emphasized that nutritionally, these wild plants could be comparable to or even sometimes superior to the

introduced cultivars.

Earlier researchers have evaluated the nutritional composition of different types of wild plants and suggested their significance as source of nutrients (Freiberger *et al.*, 1998), Markovic *et al.* (1996), Mahapatra *et al.* (2012) and Mahadkar *et al.* (2012). Recent studies on agro-pastoral societies in Africa indicated that these wild plant resources play a significant role in nutrition, food security, and income generation (Edmons and Chweya 1995). Aberoumand and Deokula (2009) have also suggested the incorporation of edible wild and semi cultivated plant resources to nutritionally marginal populations in developing countries. Variations in the nutritional content and morphological traits of fruits at inter specific level have been demonstrated in plant community (Debussche *et al.*, 1987; Herrera, 1987; and Corlett, 1996), whereas reports on intra specific variations in the quantity of nutrients particularly in the wild fruits are rare (Jordano, 1984 and Izhaki *et al.* (2002).

Bakare *et al.* (2000), Santhi *et al.* (2011) and Ullah *et al.* (2011) have evaluated the nutrient and phytochemical constituents in *M.charantia* and identified its immense value as a biomedicine. Although the general biochemical composition of immature fruits of *M.charantia* is similar to other cucurbits, bitter gourd possess comparatively high concentration of nutrients (Miniraj *et al.* 1993, Desai and Musmade, 1998 and Behera 2004).

However, considerable variations were reported in the nutrient contents in the cultivars of *M.charantia* (Wills *et al.*, 1984; Kale *et al.*, 1991; Yuwai *et al.*, 1991; Bakare *et al.* 2010 and Ullah *et al.* 2011). Studies on the nutritional constituents in *M.dioica* were reported by Singh *et al.*, (2009) and Aberoumand (2010). According to Bakare *et al.*, (2010) the future breeding and genetic emphasis in bitter gourd improvement should be placed on the development of nutritious, high yielding cultivars. But reports on the nutritional analysis in the wild varieties of *M.charantia*, a source of germplasm for crop improvement, are lacking except that of Ullah *et al.* (2011). In the present investigation, a comparative analysis of the wild and cultivated varieties of *M.charantia* was carried out to assess the intra specific variability in nutritional composition in relation to the cultivars. Correlation between the fruit morphology and the nutrient composition in the wild was also attempted to analyse the intra specific relationship.

## MATERIALS AND METHODS

Two accessions of the cultivated variety, *M.charantia* var. *charantia* (MC1 & MC2) and thirteen accessions of the wild variety *M.charantia* var. *muricata* (MC3-MC15) collected from various parts of the state of Kerala, Tamil Nadu and Karnataka were analysed in the present investigation. The characteristics of fruits and seeds of the accessions are presented in Table 1.

Table 1. Accessions of the varieties of *M. charantia* and their characteristics.

Species – accessions	Code	Collection place	Fruit shape	Nature of tubercle	Density of tubercle	Seed color	Seed shape	Extent of seed sculpture
<i>M.charantia</i> var. <i>charantia</i>	MC1	Kerala Agricultural University	Elliptical	Raised & blunt	Dense	Straw	Broad rectangular	Markedly sculptured
“	MC2	Kulathupuzha (Trivandrum)	Elliptical	Raised & blunt	Dense	Straw	Broad rectangular	Markedly sculptured
<i>M.charantia</i> var. <i>muricata</i>	MC3	Karaikal (Tamil Nadu)	Cylindrical	Raised & pointed	Sparse	Straw	Broad rectangular	Markedly sculptured
“	MC4	Aryankavu (Kollam)	Rhomboid	Raised & blunt	Sparse	Black & brown patched	Broad rectangular	Markedly sculptured
“	MC5	Achankovil (Kollam)	Globular	Raised & blunt	Medium	Straw	Broad rectangular	Markedly sculptured

Table 1. Continuation.

Species - accessions	Code	Collection Place	Fruit Shape	Nature of Tubercle	Density of Tubercle	Seed Color	Seed Shape	Extent of Seed Sculpture
<i>M.charantia</i> var. <i>muricata</i>	MC6	Kulathupuzha (Trivandrum)	Rhomboid	Raised & blunt	Medium	Black & brown patched	Broad rectangular	Markedly sculptured
“	MC7	Neyyadam (Trivandrum)	Globular	Soft & flat	Sparse	Straw	Broad rectangular	Markedly sculptured
“	MC8	Sulthan Battery (Wayanad)	Disc	Soft & flat	Sparse	Straw	Broad rectangular	Markedly sculptured
“	MC9	Pathanamthitta	Rhomboid	Raised & blunt	Medium	Straw	Broad rectangular	Markedly sculptured
“	MC10	Manipal (Karnataka)	Others	Raised & blunt	Medium	Brownish tan	Narrow rectangular	Feebly sculptured
“	MC11	Palakkad	Oblong	Raised & pointed	Sparse	Whitish brown	Narrow rectangular	Feebly sculptured
“	MC12	Kuthirankayattom (Thrissur)	Globular	Soft & flat	Sparse	White	Narrow rectangular	Feebly sculptured
“	MC13	Puthuvayal	Disc	Soft & flat	Sparse	Whitish brown	Narrow rectangular	Feebly sculptured
“	MC14	Chennai	Disc	Raised & pointed	Sparse	White	Narrow rectangular	Feebly sculptured
“	MC15	Thrissur	Cylindrical	Raised & pointed	Sparse	Whitish brown	Narrow rectangular	Feebly sculptured

### *Pomological and Nutritional Evaluation*

The accessions of the wild variety, *M.charantia* var. *muricata*, have shown variations with regard to the shape of the fruits, nature and density of tubercles, and the colour, shape and surface ornamentation of seeds. Pomological traits were evaluated using the descriptors proposed by Joseph and Antony (2010). Healthy and disinfected fresh fruits of all the accessions without seeds were used for the analysis after these were properly washed with water and dried by blotting.

Standard methods were followed for analyzing the nutrient parameters like total carbohydrate (1962), total sugar (Rangana 1979), protein (Lowry *et al* 1951) and reducing sugar (Miller 1972). The quantity of non-reducing sugar was calculated by subtracting the amount of reducing sugar from that of the total sugars. The quantity of vitamins like thiamine, riboflavin, tocopherol (Okwu, 2005) and ascorbic acid (Harris and Ray 1935) was determined. Elemental analysis included the determination of Na, Ca, Mg, Fe, Mn, Zn, and Cu. For element quantification, 0.5 g of fine powdered sample of each fruit was digested following wet digestion procedure using conc. HNO<sub>3</sub> and HClO<sub>3</sub> and filtered through Whatmann 42 filter paper and the filtrate was analysed using Atomic Absorption Spectrophotometer (Vogel, 1962).

### *Statistical Analysis*

One Way ANOVA was used to determine the significant variations in the nutritional composition between the accessions. The pomological traits and nutrient composition in the wild and cultivated varieties of *M.charantia* were subjected to PCA and Pearson Correlation study to determine the relationships between nutrient composition and pomological traits. Fruit morphological traits such as fruit weight, fruit diameter, fruit length and clutch size (number of seeds / fruit) were considered in the correlation analysis. Principal Component Analysis was carried out to detect the most variable nutrient parameters. Cluster analysis of the preliminary nutritional data was also performed using the Unweighted Pair group Method with Arithmetic Averages (UPGMA) to examine the similarity and dissimilarity at the intra specific level using NTSYS software package.

## RESULTS AND DISCUSSION

*Pomological and Nutritional Evaluation*

The wild accessions differed markedly in the shape of the fruits and seeds, nature and density of tubercles on the fruit surface and in the seed surface characteristics (Table 1). However, no variation was observed in the cultivated variety. Analysis of the nutrient composition in the accessions of the wild and cultivated varieties of *M.charantia* (Table 2 & Fig. 1) revealed higher amount of carbohydrate, reducing sugar, non-reducing sugar, protein and vitamin content in the wild varieties of *M.charantia* compared to that of the cultivated variety. However, the commercially released cultivated variety MC1 exhibited very low percentage of protein, total sugar, reducing sugar, vitamin B1, Ca, and Fe. Out of the 13 wild accessions, MC12 from Thrissur district, Kerala had the highest concentration of Vit. E, B2, Cu and Ca. The wild accession MC10 had higher concentration of carbohydrate, non-reducing sugar and Vit. B1. The accessions MC2, MC7, MC15 were characterized by higher concentration of protein, of Vit. C and reducing sugar, respectively.

The quantity of carbohydrate, reducing sugar, non-reducing sugar, protein and vitamin content in the fruits of the cultivated and wild varieties of *M.charantia* (MC1 - MC15) shown in Table 2 revealed that the accessions of the wild variety were superior in nutritional composition than the cultivated ones. However, significant variations were observed among the varieties in the content of the nutrients present in them. Data on the analysis of elements including Cu, Ca, Fe, Mn, Mg, Na, and Zn are shown (Fig.1). Analysis of the nutrient composition (Table 2 & Fig.1) revealed higher amount of carbohydrate, reducing sugar, non-reducing sugar, protein and vitamin content in the wild varieties of *M.charantia* compared to that of the cultivated variety. However, the commercially released cultivated variety MC1, exhibited very low percentage of protein, total sugar, reducing sugar, vitamin B1, Ca, and Fe. Out of the 13 wild accessions, MC12 from Thrissur district, Kerala had the highest concentration of Vit. E, B2, Cu, and Ca. The wild accession MC10 had higher concentration of carbohydrate, non-reducing sugar and Vit. B1. The accessions MC2, MC7, MC15 were characterized by higher concentration of protein, of Vit. C, and reducing sugar respectively. The micronutrients were appreciably high in the wild accessions especially in MC13, MC14, and MC8. Ullah *et al.* (2011) recorded the highest amount of total sugar, proteins, and Cu in the wild variety *muricata* from Bangladesh. Better nutritional value of some wild fruits compared to the cultivated ones was identified by Eromosele (1991) and Murugkar and Subbulakshmi (2005).




Fig. 1. Elemental Analysis in the Accessions of *M.charantia* (mg/100gm)  
 MC 1& MC2 – *M.charantia* var. *charantia* accessions  
 Mc3 MC15 – *M.charantia* var. *muricata* accessions

Table 2. Nutritional Constituents of the Cultivated and Wild Varieties of *M. charantia* (mg/1gm) - Mean  $\pm$  Standard error.

Plant	Carbohydrate	Protein	Total sugar	Reducing sugar	Non-reducing sugar	Vit. E	Vit. C	Vit. B1	Vit. B2
MC1	0.88 $\pm$ 0.003	0.32 $\pm$ 0.01	0.26 $\pm$ 0.003	0.023 $\pm$ 0.00	0.24 $\pm$ 0.008	0.23 $\pm$ 0.003	0.4 $\pm$ 0.006	0.006 $\pm$ 0.00	0.08 $\pm$ 0.006
MC2	0.84 $\pm$ 0.02	0.324 $\pm$ 0.003	0.264 $\pm$ 0.009	0.043 $\pm$ 0.006	0.22 $\pm$ 0.04	0.14 $\pm$ 0.003	0.02 $\pm$ 0.01	0.006 $\pm$ 0.000	0.05 $\pm$ 0.009
MC3	1.98 $\pm$ 0.02	0.94 $\pm$ 0.03	0.27 $\pm$ 0.04	0.16 $\pm$ 0.008	0.11 $\pm$ 0.006	0.39 $\pm$ 0.04	0.11 $\pm$ 0.01	0.07 $\pm$ 0.02	0.94 $\pm$ 0.03
MC4	1.79 $\pm$ 0.18	0.34 $\pm$ 0.006	0.41 $\pm$ 0.02	0.13 $\pm$ 0.03	0.28 $\pm$ 0.03	0.63 $\pm$ 0.04	0.02 $\pm$ 0.007	0.104 $\pm$ 0.01	0.59 $\pm$ 0.03
MC5	1.96 $\pm$ 0.04	1.26 $\pm$ 0.21	0.304 $\pm$ 0.03	0.12 $\pm$ 0.003	0.18 $\pm$ 0.02	0.71 $\pm$ 0.05	0.39 $\pm$ 0.015	0.08 $\pm$ 0.012	0.62 $\pm$ 0.05
MC6	1.94 $\pm$ 0.05	1.59 $\pm$ 0.14	0.29 $\pm$ 0.05	0.14 $\pm$ 0.006	0.15 $\pm$ 0.03	0.51 $\pm$ 0.06	0.29 $\pm$ 0.02	0.75 $\pm$ 0.04	1.31 $\pm$ 0.17
MC7	1.79 $\pm$ 0.06	1.53 $\pm$ 0.07	0.314 $\pm$ 0.01	0.15 $\pm$ 0.03	0.164 $\pm$ 0.05	0.84 $\pm$ 0.12	0.69 $\pm$ 0.18	0.11 $\pm$ 0.02	0.68 $\pm$ 0.02
MC8	1.03 $\pm$ 0.06	0.91 $\pm$ 0.06	0.31 $\pm$ 0.02	0.13 $\pm$ 0.02	0.18 $\pm$ 0.03	0.79 $\pm$ 0.12	0.399 $\pm$ 0.06	0.05 $\pm$ 0.02	0.93 $\pm$ 0.05
MC9	2.05 $\pm$ 0.06	1.63 $\pm$ 0.16	0.29 $\pm$ 0.02	0.081 $\pm$ 0.003	0.2 $\pm$ 0.02	0.65 $\pm$ 0.05	0.29 $\pm$ 0.06	0.75 $\pm$ 0.02	1.03 $\pm$ 0.02
MC10	1.29 $\pm$ 0.03	0.74 $\pm$ 0.01	0.28 $\pm$ 0.02	0.13 $\pm$ 0.02	0.89 $\pm$ 0.02	0.11 $\pm$ 0.02	0.09 $\pm$ 0.01	2.06 $\pm$ 0.15	0.005 $\pm$ 0.00
MC11	2.09 $\pm$ 0.07	1.17 $\pm$ 0.08	0.27 $\pm$ 0.02	0.09 $\pm$ 0.02	0.18 $\pm$ 0.02	0.52 $\pm$ 0.08	0.284 $\pm$ 0.02	1.36 $\pm$ 0.05	0.96 $\pm$ 0.06
MC12	0.66 $\pm$ 0.06	1.51 $\pm$ 0.14	0.32 $\pm$ 0.02	0.05 $\pm$ 0.02	0.27 $\pm$ 0.04	0.96 $\pm$ 0.03	0.07 $\pm$ 0.02	0.12 $\pm$ 0.02	1.52 $\pm$ 0.095
MC13	1.63 $\pm$ 0.003	1.28 $\pm$ 0.03	0.31 $\pm$ 0.03	0.05 $\pm$ 0.01	0.26 $\pm$ 0.02	0.87 $\pm$ 0.09	0.25 $\pm$ 0.01	1.84 $\pm$ 0.11	0.85 $\pm$ 0.07
MC14	0.59 $\pm$ 0.02	1.48 $\pm$ 0.11	0.3 $\pm$ 0.04	0.14 $\pm$ 0.02	0.16 $\pm$ 0.02	0.83 $\pm$ 0.05	0.27 $\pm$ 0.04	1.78 $\pm$ 0.07	0.98 $\pm$ 0.04
MC15	2 $\pm$ 0.06	1.3 $\pm$ 0.04	0.28 $\pm$ 0.02	0.19 $\pm$ 0.04	0.09 $\pm$ 0.02	0.42 $\pm$ 0.04	0.19 $\pm$ 0.03	1.97 $\pm$ 0.04	0.64 $\pm$ 0.06
F value	78.2***	24.04***	2.00NS	7.68***	49.53***	18.69***	10.86***	216.74***	49.95***

MC 1& MC2 - *M.charantia*var.*charantia*accessions MC3-MC15M.*charantia*var.*muricata*accessions

The present study revealed that the accession MC7, with considerably rich vitamin C content (69 mg/100gm), can be used in the bitter gourd breeding programme. According to Dey *et al.* (2005-2006), breeding for nutritional/medicinal quality of bitter gourd typically emphasizes accessions with relatively high Vit. C content. The present study does not recommend the selection of MC2 with very low Vit.C. content. The study also revealed the superiority of the accession MC12, and it recommends the variety for hybridization programme. Decrease in the nutritional quality in the cultivated variety, *M. charantia* var. *charantia*, may be due to the inappropriate method of selection during domestication, where importance was given to those plants having larger plant parts like larger fruits. The study is not in agreement with the view proposed by Kendrick *et al.* (2004) that the nutritional quality of *M.charantia* decreased as a result of domestication.

### Statistical Analysis

Pearson correlation study of fruit morphological traits and nutritional parameters of the accessions of the wild variety, *M.charantia* var. *muricata*, revealed significant negative correlation between the nutrients like protein, Vit. C, Vit. E, and Vit. B2 and fruit characteristics like fruit weight, diameter, length and clutch size, i.e., number of seeds / fruit at 0.05 level (Table 3a & 3b). However, a positive correlation was observed between fruit length and clutch size. Negative correlation was also existed between the quantity of minerals like Cu, Ca, and different fruit characters. It was observed that the carbohydrate content and fruit traits were positively correlated while reducing sugar showed significant positive correlation only with fruit size. Positive correlation was also noted between total sugar and weight and diameter of the fruits and not with fruit length. No significant correlation was observed between fruit length and quantity of Fe. Positive and negative correlation also existed between some of the nutritional parameters (eg: Mg, Zn, Fe, Na) with the fruit traits at 0.01 and 0.05 levels. Principal Component Analysis (Table 4) revealed that the quantity of carbohydrate, Mg, Na, and protein content are the highly loaded characters in the first two principal axis with the percentage of variance 65.59 and 15.62, respectively. Variations were observed at the intra specific levels in *M.charantia* var. *muricata* during the investigation which may be due to the influence of those characters.

Table 3a. Correlation Between Pomological and Nutritional Value of *M.charantia* var. *muricata* Accessions.

	Na	No: seeds	NR. Sugar	PTN	RD. Sugar	T.Sugar	Vit B1	Vit B2	Vit E	Zn
Na	1									
No: seeds	-0.14	1								
NR. Sugar	-0.35*	0.24*	1							
PTN	-0.05	-0.56**	-0.27**	1						
RD. Sugar	0.17	0.13	0.29**	0.21*	1					
T.Sugar	-0.17	0.02	-0.18	-0.47	-0.14	1				
Vit B1	-0.59**	0.15	0.62**	0.09	0.20	-0.41**	1			
Vit B2	0.15	-0.42**	-0.63**	0.57**	0.389**	-0.05	-0.34**	1		
Vit E	-0.15	0.57**	-0.62**	0.43**	-0.48**	0.39**	-0.37**	0.59**	1	
Zn	0.33**	0.23*	0.08	-0.44**	0.25*	0.09	-0.37**	-0.396**	-0.25*	1

\*Correlation is significant at the 0.05 level (2-tailed).

\*\* Correlation is significant at the 0.01 level (2-tailed).

F.L. - Fruit Length, F.W. - Fruit Weight, F.D. - Fruit diameter, N.R.-sugar - Non-reducing sugar, PTN-Protein, R.D.sugar - Reducing sugar, T.sugar - Total sugar

Table 3b. Correlation Between Pomological and Nutritional Value of *M.charantia* var. *muricata* Accessions.

	Vit C	Ca	CHO	Cu	Fe	F.W	F.D	F.L	Mg	Mn
Vit C	1									
Ca	-0.28**	1								
CHO	-0.07	-0.42**	1							
Cu	-0.17	0.88**	-0.44**	1						
Fe	0.04	0.198*	-0.22*	-0.05	1					
F.W	-0.47**	-0.33**	0.53**	-0.33**	-0.34**	1				
F.D	-0.37**	-0.30**	0.47**	-0.27**	-0.32**	0.77**	1			
F.L	-0.51**	-0.36**	0.46**	-0.49**	-0.01	0.51**	0.41**	1		
Mg	-0.08	0.25*	-0.54**	0.19	-0.58**	-0.52**	-0.47**	-0.13	1	
Mn	0.12	0.37**	-0.7**	0.25*	0.79**	-0.60**	-0.46**	-0.26**	0.72**	1

\*Correlation is significant at the 0.05 level (2-tailed). \*\* Correlation is significant at the 0.01 level (2-tailed).

FL-Fruit Length, F.W-Fruit Weight ,F.D- Fruit diameter

Table 4. Highly Loaded Characters in Principal Component Analysis.

Characters	Principal Axis 1	Principal Axis 2
CHO	1.504	0.070
Mg	1.497	-0.788
Na	1.292	1.101
Protein	0.645	-0.107
Eigen values	9.839	2.344
Percentage	65.593	15.624
Cumulative percentage	65.593	81.217

The dendrogram constructed based on the nutrient composition (Fig. 2) revealed two principal clusters at an Euclidean distance of 0.68. Each principal cluster had two subclusters with euclidean distances 0.62, 0.68 for the first and 0.40 and 0.52 for the second subcluster, respectively. The wild accessions MC13, MC14, and MC15 were clustered in the second subcluster at an euclidean distance of 0.5. The dendrogram did not differentiate the wild and cultivated taxa. Clustering of the wild and cultivated accessions together in the first principal cluster suggest that they possessed more or less the same type of nutritional components. However, the accessions differed in the quantity of nutrients. It was observed that the cluster distance between the wild accessions MC8 and MC5, MC 6 and MC12, MC 7 and MC 3, and MC14 and MC 13 was minimum based on nutritional value. The grouping of MC13 and MC 14 together in the second cluster may be due to the presence of high quantity of micronutrients in those accessions. Similarity based on the nutrient characters was evident from the grouping of the wild accessions MC8 and MC5 (0.22) , MC6 and MC12 (0.28), MC7 and MC3 (0.30) and the two cultivated accessions MC1 and MC2 (0.27) The least cluster distances observed in those cited above revealed that they are closely related based on the nutrient characters. The study noticed (Table 2) that among the cultivated variety, the cultivar Preethi , procured from Kerala Agricultural University had comparatively more nutrient contents than the land race collected from Kulathupuzha and that may be due to the appropriate agricultural practices employed to the cultivated species.

UPGMA analysis revealed that the clustering of wild accessions was not in accordance with fruit and seed characters, i.e., the accessions with markedly sculptured seeds (MC3-MC9) and feebly sculptured seeds(MC10- MC15) were grouped intermittently. It indicates that the seed characters have no role in determining the nutritional quality. However, clustering of the three accessions MC13,14&15 together in a group revealed the interrelationship of the taxa. Scatter analysis based on nutrient characters (Fig. 3) also supported the grouping of wild accessions and the two cultivated accessions as in the



UPGMA dendrogram. Occurrence of wild accessions in independent groups in different axes in the scatter plot points out the nutritional variation, and that may be due to genetic and environmental interaction. Obeso and Herrera (1994) suggested that the variations in nutritional characters among species might be due to the consequences of genetic and environmental factors.




Fig 2. UPGMA Dendrogram based on Nutritional Parameters.




Fig. 3. Scatter Plot Based on Nutritional Characters.

#### *Correlation between fruit traits and nutrient contents in the wild varieties of M.charantia*

The correlation studies between the nutritional characters and fruit traits will be of help in selecting the wild accessions with higher nutrient content for crop improvement programmes. Pearson correlation study among the wild varieties of *M.charantia* revealed strong negative correlation between fruit traits (fruit weight, diameter, and length) and Vit. C, copper, and calcium indicating the presence of rich nutrient contents in plants with small fruits (Table 4).



According to Behera *et al.*, (2010), negative correlation was observed between fruit traits (weight, fruit length and diameter) and ascorbic acid content in the cultivars of *M.charantia*. They suggested that the selection of small fruited cultivars is known to improve the ascorbic acid content. The present study thus recommends the accession MC7 with high vitamin C (69 mg/gm), collected from Thiruvananthapuram district Kerala as the most suitable taxa and can be used to enhance the ascorbic acid content of the cultivated variety. The negative correlation between fruit size and protein was reported at intra specific level in the fruits of *Rbhamnus alaternus* (Izhakiet al. 2002). Correlation between protein content and fruit size observed in the present findings indicates that the wild variety MC2 with small sized fruits can also be preferred for crop improvement. A positive significant correlation between all the four fruit traits and carbohydrate content noticed in the wild varieties indicates that carbohydrate is directly related to the size of the fruit. Significant negative correlation between fruit size and Ca content suggests that small fruits are rich sources of Calcium. Bakare *et al.* (2010) reported that Ca was the most abundant mineral present in the leaf of *M.charantia*. The present investigation observed a positive significant correlation between reducing and non-reducing sugar with fruit size and fruit weight, respectively. Markovik *et al.* (1996) reported a negative correlation between fruit weight and total sugar content in tomato.

The study noticed both positive and negative correlation between fruit traits and the nutritional parameters like total sugar, Vit.B<sub>1</sub> and Zn, at 0.01 and 0.05 levels indicating the antagonistic interaction between nutrients ( eg:CHO and vit E, Ca and Mg). Cecil *et al.* (1995) pointed out that the interrelationship among fruit nutrients may be due to synergistic and antagonistic interaction between nutrients or due to plant-environment relations. Significant positive correlation existing between Mg and Ca ( $r=0.25$ ) at 0.05 level suggests that they are closely associated in the metabolic pathways. Such an association was detected by earlier researchers (Garten,1976; Alonso and Herrera, 2001,Izhaki *et al.* 2002) and Garten (1976) opined that this correlation was probably a consequence of the close association of Ca and Mg in metabolism and photosynthesis.

Most of the fruit morphological traits were significantly correlated with each other at 0.01 level too. There was a direct relationship between the nutrient contents and the fruit traits (fruit weight and length together with fruit diameter and clutch size). Behera *et al.* (2010), opined that the fruit weight had the greatest direct effect on fruit length in the cultivars of bitter melon. Variations observed in the nutrient composition in the fruits of the wild accessions in the present investigation may be due to the interaction of the environment and the genotype. Difference in the environmental conditions between plant microsites impressed greater variability on fruit composition than on fruit morphological traits.

## CONCLUSION

The study revealed the superiority of the wild variety, *M.charantia* var. *muricata*, over the cultivar in terms of nutrient composition. The study recommends the use of the wild accessions, particularly MC10, MC12, MC 2 & Mc7, the rich sources of nutrients to produce varieties with superior qualities in *Momordica* breeding programmes. The investigation also emphasized the conservation of the wild varieties for future breeding programme.

## ACKNOWLEDGEMENT

The authors acknowledge the Department of Science and Technology for the financial support and Dr. P.M. Radhamany, Head, Department of Botany, University of Kerala, for providing the facilities.

## REFERENCES

- ABEROUMAND, A. (2010) A comparative study of nutrients and mineral molar ratio of some plant foods with recommended dietary allowances. *Advanced Journal of Food Science and Technology* 2 (2): 104-108.
- ABEROUMAND, A. and S.S. DEOKALA. 2009. Proximate and mineral composition of wild coco (*Eulophia create* L.) tubers in Iran. *Asian Journal of Food Agriculture and Industry* 2(02) 203-209.
- ALONSO, C. and C.M. HERRERA. 2001. Patterns made of patterns: variation and covariation of leaf nutrient concentrations within and between populations of *Prunus mabaleb*. *NewPhytologist* 150 629-640.
- BAKARE, R.I., O.A. MAGBAGBEOLA, A.I. Akinwande, and O.W. Okunowo. 2010. Nutritional and chemical evaluation of *Momordica charantia*. *Journal of Medicinal Plant Research* 4(21) : 2189-2193.
- BEHERA, T. K. 2004. Heterosis in bitter gourd. In *Hybrid vegetable development*. eds. P. K. Singh, S. H. Dasgupta and S. K. Tripathi. Haworth Press, New York.
- BEHERA, T. K., S. BEHERA, L.K. BHARATHI, K. JOSEPH JOHN, W.S. PHILIPP, and E.S. JOHN. 2010. Bitter gourd: Botany, horticulture, breeding. *Horticultural Reviews* 37: 101-141.
- BHARATHI, L.K., A.D. MUNSHI, T.K. BEHERA, VINOD, K. JOSEPH JOHN, A.B. DAS, K. V. BHAT, and A.S. SIDHU. 2012. Production and preliminary characterization of inter-specific hybrids derived from *Momordica* species. *Current Science* 103(2):178-186.
- CECIL, J.S., G.E. BARTH, N.A. MAIER, W.L. CHVYL, and M. N. BARTESKO. 1995. Leaf chemical composition and nutrient removal by stem of *Leucadendron* cvSilvan Red and Safari Sunset. *Australian Journal of Experimental Agriculture* 35: 275-283.
- CORLETT, R. T. 1996. Characteristics of vertebrate- dispersed fruits in Hong Kong. *Journal of Tropical Ecology* 12: 819-833.
- DEBUSSCHE, M., J. CORTEZ, and I. RIMBAULT. 1987. Variation in fleshy fruit composition in the Mediterranean region: the importance of ripening season, life form, fruit type and geographical distribution. *Oikos* 49: 244-252.
- DESAI, U. T., and A.M. MUSMADE. 1998. Pumpkins, squashes and gourds 273-298. In *Handbook of vegetable science and technology: production, composition, storage and processing*. eds. D. K. Salunkhe and S. S. Kadam Marcel Dekker, New York.
- DEY, S.S., T.K. BEHERA, and C. KAUR. (2005-2006). Genetic variability in ascorbic acid and carotenoids content in Indian bitter gourd (*Momordica charantia*L.) germplasm. Cucurbit Genetics Cooperative Report 28-29: 91-93.
- EDMONS, J., and J. CHWEYA. 1995. Black nightshades, *Solanum nigrum* L. and related species in Promoting the conservation and use of under utilized and neglected crops. 221-234. Taylor and Francis London.
- EROMOSELE, I.C., C.O. EROMOSELE, and M. KUZHKUZHA. 1991. Evaluation of mineral elements and ascorbic acid contents in fruits of some wild plants. *Plant Foods for Human Nutrition*.(41)151-154.
- FREIBERGER, C.E., D.J. VANDERJAGT, A. PASTUSZYN, R.S. GLEW, G. MOUNKAILA, M. MILLSON, and R.H. GLEW. 1998. Nutrient content of the edible leaves of seven wild plants from Niger. *Plant Foods for Human Nutrition*. 53(1):57-69.
- GARTEN, C.T. 1976. Correlation between concentration of elements in Plants. *Nature* 261: 686-688.
- HARRIS, L. J., and S.N. RAY. 1935. Determination of plasma ascorbic acid by 2,6-dichlorophenol indophenol titration. *Lancet* 1:462.
- HEDGE, J.E., and B.T. HOFRIETER. 1962. In: Carbohydrate chemistry 17. eds. R.L. Whistler and J. N. Be Miller. Academic Press, New York.

- HERRERA, C. M. 1987. Vertebrate- dispersed plants of Iberian Peninsula: a study of fruit characteristics. *Ecological Monographs* 57: 305-331.
- IZHAKI, I., E. TSAHAR, I. PAULY, and J. FRIEDMAN. 2002. Within population variation and interrelationship between morphology, nutritional content and secondary compounds of *Rhamnus alaternus* fruits. *New Phytologist* 156: 217-223
- JORDANO, P. 1984. Seed weight variation and differential avian dispersal in blackberries *Rubus ulmifolius*. *Oikos* 43:149-153.
- JOSEPH J.K., and V.T.ANTONY. 2010. A taxonomic revision of the genus *Momordica* L. (Cucurbitaceae) in India. *Ind. J. Plant Genet. Res.* 23(2): 172-184.
- KALE, A.A., S.R. GADAKH, and R.N. ADSULE. 1991. Physico-chemical characteristics of improved varieties of bittergourd (*Momordica charantia*L.) *Maharashtra Journal of Horticulture* 5:56-59.
- KENDRICK, L. M., X.Y. MEI, and N.K. BHATTARAI. 2004. Allozyme, morphological and nutritional analysis bearing on the domestication of *Momordica charantia*L. (Cucurbitaceae). *Economic Botany* 58(3): 435-455.
- LOWERY, O.H., N.J. ROSEBROUGH, L.A. FAN, and R.J. RANDALL. 1951. Protein measurement with the Folin-Phenol reagent. *Journal of Biological Chemistry* 193:262-275.
- MAHADKAR, S., S. VALVI, and V. RATHOD. 2012. Nutritional assessment of some selected wild edible plants as a good source of mineral. *Asian Journal of Plant Science Research* 2 (4):468-472.
- MAHAPATRA, A. K., S. MISHRA, U.C. BASAK, and P.C. PANDA. 2012. Nutrient analysis of some selected wild edible fruits of deciduous forests of India: an explorative study towards non conventional bio-nutrition. *Advanced Journal of Food Science and Technology* 4(1): 15-21.
- MARKOVIC, Z., J. ZDRAVKOVIC, and M. DAMJANOVIC. 1996. Correlation between the morphological characteristics and the biochemical components of tomato fruit quality. *Acta Horticulture* (ISHS) 462: 151-156.
- MILLER, G. I. 1972. Use of dinitrosalicylic acid reagent for determination of reducing sugars. *Analytical Chemistry* 31:426.
- MINIRAJ, N., K.P. PRASANNA, and K.V. PETER. 1993. Bitter gourd *Momordica spp.* In Genetic improvement of vegetable plants. eds. G.Kaloo and B.O. Bergh. Pergamon Press, Oxford, UK.
- MURUGAR, D. A., and G. SUBBULAKSHMI. 2005. Nutritive values of wild edible fruits, berries, nuts, roots and spices consumed by the Khasi tribes of India. *Ecology and Food Nutrition* 44 (3): 207-223.
- OBESO, J.R. and C.M. HERRERA. 1994. Inter and intra specific variation in fruit traits in co occurring vertebrate dispersed plants. *International Journal of Plant Science* 155: 382-387.
- OKWU, D. E. 2005. Phytochemicals, vitamins, and mineral contents of two Nigerian medicinal plants. *International Journal of Molecular Medicine and Advanced Science* 1(4): 375-381.
- RANGANA, S.C. 1979. Manual of Analysis of Fruit and Vegetable Products. Tata McGraw Hill Publishing Company Limited, New Delhi.
- REYES, M.E.C., B.H. GILDEMACHER, and G.J. JANSEN. 1994. *Momordica L.*, pp.206-210 In: *Plant resources of South-East Asia: Vegetables*. eds. J.S. Siemonsoma and K. Piluekaeds. Pudoc scientific publishers, Wageningen, the Netherlands.
- SINGH, D., V. BAHADUR, D.B. SINGH, and G. GHOSH. 2009. Spine gourd (*Momordica dioca*) : An underutilized vegetable with high nutritional and medicinal values. *Acta Horticulturae* (ISHS) 809: 241-248.
- ULLAH, M., F. KARIMCHY, S.K. SARKAR, M.K. ISLAM, and N. ABSAR. 2011. Nutrient and phytochemical analysis of four varieties of bitter gourd (*Momordica charantia*) grown in Chittagong Hill Tracts, Bangladesh. *Asian Journal of Agricultural Research*: 1-5.

- VOGEL, A.I. 1962. In: A text-book of quantitative inorganic analysis (including elementary instrumental analysis). 3rd Ed. Longmans, Green, London.
- WILLS, R.B.H., A.W.K WONG, and F.M. SCRIVEN. Greenfield, H.1984. Nutrient composition of Chinese vegetables. *Journal of Agriculture and Food Chemistry*.32(2):413-416.
- YUWAI, K.E., K.S. RAO, J.C. KALUWIN, G.P. JONES, and D.E. RIVETTS. 1991. Chemical composition of *Momordica charantia*L. Fruits. *Journal of Agriculture and Food Chemistry* 39: 1782-1763.