

Role of Secondary Metabolites and Radical Scavenging Aptitude for Better Adaptability of Mangroves in Varying Salinity of Sundarbans, India

Nirjhar Dasgupta^{1,2}, Chandan Sengupta² and Sauren Das¹

¹Agricultural and Ecological Research Unit, Indian Statistical Institute
203, B.T.Road, Kolkata 700108, India, ²Department of Botany, University of Kalyani
Kalyani, Nadia, 741235, India

ABSTRACT

Comparative adaptability in five halophytes (*Bruguiera gymnorrhiza*, *Excoecaria agallocha*, *Heritiera fomes*, *Phoenix paludosa* and *Xylocarpus granatum*, of which, *H. fomes* and *X. granatum* presently are stressed in Sundarbans area) were evaluated with respect to occurrence of total phenol, flavonoids, and radical scavenging ability following ABTS [2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)], DPPH (1, 1-diphenyl-2-picrylhydrazyl) assay and Fe₂⁺ chelating ability.

Amount of total phenol (TP) and flavonoids (TF) were much higher in all taxa grown in Sundarbans than those of mesophytic one. TP and TF were significantly augmented as the substrate salinity increased in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* but disordered in *H. fomes* and *X. granatum*, where increment occurred only up to a certain salinity level. Percent of free radical scavenging of extractants by DPPH and ABTS radical perceived significant correlation with salinity in former three but differ in *H. fomes* and *X. granatum*. Ferrous ion chelating ability also showed the similar trend.

Owing to polyphenols occurrence and ROS scavengers, the present work clearly indicates the better adaptability of *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* in elevated substrate salinity than those of the other two. Lower ROS scavenging ability of *H. fomes* and *X. granatum* also points to their perilous occurrence in elevated saline zones.

Key words: Flavonoids; Phenols; ABTS; DPPH; Fe₂⁺ chelation; mangroves; Sundarbans.

Correspondence : S. Das Address: Agricultural and Ecological Research Unit, Indian Statistical Institute
203, B.T.Road, Kolkata 700108, India E-mail: sauren@isical.ac.in

DOI: 10.32945/atr3621.2014

INTRODUCTION

Mangroves considered as front line defense of tropical and subtropical coastal and estuarine ecosystem, can efficiently mitigate sea surges (McIvor *et al.*, 2012). But unfortunately, world-wide mangrove area deteriorated from 198,000 km² to 146,530 km² since 1980 to 2000 (FAO, 2007) that while a more recent estimation described as 137,760 km² which is about 0.7% of the total tropical forest of the world (Giri *et al.*, 2011). These indicate the magnitude of mangrove loss, and calls for serious attention to the significance of mangrove restoration programme. The mangrove formation has immense importance to the coastal inhabitants in both productive (resources of forest products) and protective ways (against cyclonic violence, soil erosion etc.) (Alongi, 2009).

In the Indian subcontinent, Sundarbans mangrove swamp is the world's largest and having richest floral diversity (21°31' – 22° 30' N and 88°10' – 89°51' E), formed by two major rivers - the Ganga and the Brahmaputra and with their numerous tributaries (Dasgupta *et al.*, 2012). Environmental (tectonic upliftment in north-western part, India and subsidence in the east, Bangladesh) and demographic compulsion (damming on river, urban sewage and industrial wastes expedite siltation in the riverbed) are interrupting the fresh water discharge through river systems, leading to salinity elevation of this zone. The present soil salinity registers between 15 and 27 ppt (Nandy *et al.*, 2007). Haq *et al.* (1999) estimated that salinity has elevated by 20% in the Sunderbans since 1990. Sea level rise, human intervention and reformed ecology are the prime factors of salinity escalation which pose the major threat to the vegetation and triggered rapid annihilation to the forest and even caused extinction to some existing species (Parani *et al.*, 2000). These demographic and environmental adversities have been proved catastrophic for some important plant species of Sundarbans including *Aegialitis rotundifolia*, *Heritiera fomes*, *Nypa fruticans*, *Xylocarpus granatum* and *X. mekongensis* (Banerjee, 1999; Upadhaya *et al.*, 2002).

Considerable scientific evidences suggest that under oxidative stress, reactive oxygen species (ROS) such as superoxide, hydroxyl and peroxy radicals are generated and the stability between anti-oxidation and oxidation is supposed to be a critical concept for maintaining a biological homeostasis (Davies, 2000). Consequently, research on antioxidant principles of plants has been fast-tracked and halophytes

have been recognized for marked antioxidant potential activities (Tiwari, 2001). Polyphenols are antioxidants, having redox properties, act as reducing agents, hydrogen donors and singlet oxygen quenchers. A number of secondary metabolites were traced from mangroves periodically and have been recognized as pivotal bioactivities towards efficient slat management (Bandarnayeke, 2002; Ravikumar *et al.*, 2010, 2011). Despite of innumerable bioactivities, polyphenols perhaps are mostly attributed to antioxidant property (Sahidi, 2000; Cushnie and Lamb, 2005; Alam, 2011). Some essential phenols and flavonoids were traced from several mangroves (Krishnamoorthy *et al.*, 2011; Yongxin *et al.*, 2012). A variety of antioxidants and their free radical scavenging ability were evaluated from Indian mangroves by Vadlapudi and Naidu (2009).

Polyphenols in plants are well established as a defensive force against detrimental impact of abiotic stress which leads to generate free radicals (Ostrowskaa *et al.*, 2001; Zurina *et al.*, 2010; Lee *et al.*, 2012). The commonly utilized methods of antioxidant activity are ABTS [2, 2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid)] and DPPH (1, 1-diphenyl-2-picrylhydrazyl) assay. Both of them have intense reproducible characteristics under certain assay conditions. ABTS⁺ can be dissolved in aqueous and organic media, in which the antioxidant activity can be measured, due to the hydrophilic and lipophilic nature of the compounds occurring in the plant cells (Arnao, 2000). In contrast, DPPH has been used extensively as a free radical evaluating reducing substances (Cotelle *et al.*, 1996) and for studying the free radical scavenging activities of polyphenol compounds (Duan *et al.*, 2006). DPPH can only be dissolved in organic media, especially in ethanol, and this has a limitation when interpreting the role of hydrophilic antioxidants. Both radicals show similar bi-phase kinetic reactions with many antioxidants. Hydrogen peroxide itself is not very reactive, but it would provide highly reactive species OH[•] through Fenton reaction (Halliwell and Cross, 1994). Earlier reports suggested that H₂O₂ could induce DNA breakage in the intact cell (Imlay, 2003). Thus removal of H₂O₂ is decisive for cellular homeostasis. The reducing capacity of a compound may serve as indicator of its potential antioxidant ability (Meir *et al.*, 1995). The antioxidant capacity of compounds has been attributed to various factors such as prevention of chain reaction, chelating metals, reductive capacity and radical scavenging ability (Diplock, 1997; Yildirim *et al.*, 2001).

In view of this, the present work deals with five mangroves taxa of which, two are presently in emaciated occurrence in Sundarbans area,

taken from five islands with increasing soil salinity. A comparative account of the prevalence of total phenols and flavonoids among the taxa would be indicative regarding the adaptive ability in *in-situ* condition. The differential magnitude of antioxidative capability among these species has been determined by free radical scavenging result using ABTS and DPPH assay. The antioxidant ability has also been looked into by Fe_2^+ chelating capacity. All these information might be informative towards the sustainability of the species. For the sake of better appreciation of the process, the same replicas of the taxa are also taken from the mesophytic habitat.

MATERIALS AND METHODS

Leaves of well matured (about 12-15 years) five taxa were collected in which, two were in distressed condition (*Heritiera fomes* and *Xylocarpus granatum*) while the other three were profusely occurring (*Bruguiera gymnorrhiza*, *Excoecaria agallocha* and *Phoenix paludosa*) in the same regime. The collection sites were chosen from naturally occurring in five different islands according to increasing salinity gradient, namely Sajnakhali (22° 08' 48" N, 88° 50' 10" E), Sudhanyakhali (22° 07' 35" N, 88° 48' 19" E), Sonargaon (22° 06' 19" N, 88° 45' 47" E), Jharkhali (22° 02' 25" N, 88° 42' 29" E) and Dobanki (21° 58' 17" N, 88° 45' 57" E) (Fig. 1A - D). The soil salinities of these places were 11.77 ± 2.1 , 12.25 ± 1.96 , 12.4 ± 1.18 , 13.98 ± 2.29 and 15.23 ± 2.16 ppt respectively (Dasgupta *et al.* 2012). A controlled mesophytic (at the premises of Indian Statistical Institute, Kolkata where the soil salinity is 2 ppt, (Nandy *et al.*, 2009)) replicas of the studied taxa of well matured up to 12 – 15 years old, were also taken for comparative study (Table 1).

Preparation of extraction

Methanolic extracts were prepared by maceration of mature leaves (10gm) with 250ml of 95% ethanol, at room temperature for 24 hours and then filtered. The procedure was repeated thrice. The extracts were filtered and centrifuged (4500rpm, 15min). The solvent was evaporated on rotary evaporator (Neocool Circulator, CF 300, Yamato). The concentrated methanolic leaf extracts were kept frozen (-20°C) until further use (Huda-Faujan *et al.*, 2009). Leave samples were collected from three individuals of each species from each site.


**Fig. 1.**

Fig. 1. A-D. Maps of location areas. **A.** Map of India; **B.** Map of West Bengal; **C.** Map of lower Bengal basin, where the Sundarbans mangrove forest included; **D.** Expanded map of study site of the islands of Sundarbans, indicating the geographic position.

Table 1. Location and soil salinity of the collecting sites.

Name of Islands	Lat/Long values	Salinity (ppt)	Named as
ISI Garden (mesophytic)	22° 38' 43" N, 88° 22' 33" E	2.0 ± 1.21	Site 1
Sajnakhali (Sundarbans)	22° 08' 48" N, 88° 50' 10" E	11.77 ± 2.1	Site 2
Sudhanyakhali (Sundarbans)	22° 07' 35" N, 88° 48' 19" E	12.25 ± 1.96	Site 3
Sonargaon (Sundarbans)	22° 06' 19" N, 88° 45' 47" E	12.4 ± 1.18	Site 4
Jharkhali (Sundarbans)	22° 02' 25" N, 88° 42' 29" E	13.98 ± 2.29	Site 5
Dobanki (Subdarbans)	21° 58' 17" N, 88° 45' 57" E	15.23 ± 2.16	Site 6

Determination of Total Phenol

Total phenol contents in the extracts were determined following modified Folin-Ciocalteu method (Singleton and Rossi, 1965). Aliquots (0.05-0.3 mg ml⁻¹) of the extractants were prepared for all the samples. To 0.1 ml of the methanolic solution of the extractants, 2 ml of Folin-Ciocalteu reagent (previously diluted with water at 1:10 v/v) and 1.6 ml (7.5%) of Sodium carbonate (Na₂CO₃) were added. The mixture was allowed to react at room temperature for 30min for colour development. Spectrophotometric absorbance (Helios γ, Thermo Electron Corporation) was measured at 765 nm. Gallic acid was used as a positive control in the same concentrations. The total phenolic content was expressed as gallic acid equivalents (GAE) in mg g⁻¹ of dry material based on the calibration curve, where X was the absorbance and Y was the gallic acid equivalent (mg g⁻¹). Each experiment was carried out in triplicates.

Determination of Total Flavonoids

Estimation of total flavonoids in the plant extracts was carried out following Jia *et al.* (1999). Aliquots (0.2-1.2 mg ml⁻¹) of the extractants were prepared for all the samples. To 0.1 ml of methanolic solution of the extractants, 0.3 ml of 5% Sodium nitrite (NaNO₂) was added and mixed well. The mixture was kept at 25°C temperature for 5 min. Then, 0.3 ml of 10% Aluminum chloride (AlCl₃) was added. The tubes were then allowed to stand at room temperature for 5 min. After which, 1 mM Sodium hydroxide (NaOH) solution was added and the absorbance of the solution was measured at 510 nm. Quercetin was used in the same concentration as a

positive control. Total flavonoid content was calculated as quercetin equivalent (mg g^{-1}) using the following the calibration curve, where X was the absorbance and Y was the quercetin equivalent (QE mg g^{-1}). Each experiment was carried out in triplicates.

ABTS+ scavenging assay

To determine ABTS radical scavenging assay, the method of Re et al. (1997) was adopted. The working solution was prepared by mixing two stock solutions 7mM ABTS solution and 2.4 mM potassium per-sulfate solution, in equal quantities and allowing them to react for 12 hours at room temperature in the dark. This solution was diluted 60 times while working. Fresh ABTS solution was prepared for each assay. Plant extract (0.1ml) in methanol containing 0.005-0.03 mg ml^{-1} conc. of the extract were allowed to react with 3 ml of the ABTS solution and the absorbency was taken at 734 nm after 7 min. Butylated Hydroxy Toluene (BHT) in the same concentration as the sample was used as positive control. Each experimental data was taken in triplicates. The ABTS scavenging capacity of the extract was calculated as:

$$\text{ABTS radical scavenging activity (\%)} = \{(A_c - A_s)/A_c\} \times 100$$

Where A_c is the absorbance of the blank reaction and A_s is the absorbance in presence of the sample of the extracts. Each experiment was carried out in triplicates.

DPPH quenching assay

The free radical scavenging activity of different plant extracts was determined using the stable radical DPPH (1, 1-diphenyl-2-picrylhydrazyl) following Ribeiro *et al.* (2002). Freshly prepared DPPH solution (25 mg L^{-1}) in methanol was prepared and 3.9 ml from this was mixed with 0.1 ml of extract in methanol containing 0.05-0.3 mg ml^{-1} conc. of the extract. The absorbance was measured at 517 nm after 30 minutes. Butylated Hydroxy Toluene (BHT) in the same concentration as the sample was used as positive control. Each experimental data was taken in triplicates. The capability to scavenge the DPPH radical was calculated using the following equation:

$$\text{DPPH radical scavenging activity (\%)} = \left\{ \frac{A_c - A_t}{A_c} \right\} \times 100$$

Where A_c is the absorbance of the blank reaction and A_t is the absorbance in presence of the sample of the extracts. Each experiment was carried out in triplicates.

Fe₂⁺ chelating study

The chelating activity of the extracts for ferrous ion was measured following Haro-Vicente *et al.* (2006) with minor adjustments. The reaction was carried out in HEPES buffer (20 mM, pH 7.2). Various concentrations of the plant extracts (0–300 mg ml⁻¹) were added to 12.5 mM ferrous sulfate solution and the reaction was initiated by the addition of ferrozine (75 mM). The mixture was shaken vigorously and incubated for 20 min at room temperature. The absorbance is measured at 562 nm. Three repeats were done for each test. EDTA was used as a positive control.

$$\text{Fe}^{2+} \text{ Chelation ability (\%)} = \left\{ \frac{A_c - A_t}{A_c} \right\} \times 100$$

Where A_c is the absorbance of the blank reaction and A_t is the absorbance in presence of the sample of the extracts.

In all scavenging/chelating ability are expressed in terms of IC₅₀. It is defined as the concentration of the plant extract that's needed to scavenge/chelate 50% of the scavengers/chelating ions present were calculated by the following equation:

$$\text{IC}_{50} = \left\{ \frac{\text{Percent Inhibition}}{\text{Concentration of the sample}} \right\} \times 50$$

Pearson coefficient, (two-tailed bivariate correlation coefficient) was estimated for each parameter such as amount of incidence of total phenols and flavonoids, percent inhibition of free radical scavenging by DPPH and ABTS assay and percent ability to Fe₂⁺ chelation in respect to salinity increment were considered. The correlations among the said parameters were derived using SPSS (version 12.0) software.

RESULTS AND DISCUSSION

Estimated total phenol (TP) in all studied plants was much higher in plants grown in Sundarbans than those of the mesophytic individuals. In mesophytic

plants, highest TP was observed in *P. paludosa* (35.59 GAE mg g⁻¹) and in other taxa, ranged within 8.68 – 12.65 GAE mg g⁻¹. The amount increased consistently along the increasing soil salinity in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* but this trend was disrupted in *H. fomes* and *X. granatum*. Amount of total phenol increased up to site 4 in *H. fomes* (32.54 mg g⁻¹) and up to site 3 in *X. granatum* (38.48 mg g⁻¹), beyond which it decreased. TP increment in *H. fomes* was nominal from site 3 to site 4 (31.54 and 32.45 mg g⁻¹ respectively) and in *X. granatum* TP increment over mesophytic condition was up to site 3 and it went down in site 4, 5 and 6 (35.59, 31.56 and 31.48 mg g⁻¹ respectively) (Fig.2A).

The incidence of Total Flavonoids (TF) was expressed in Quercetin equivalent. In *B. gymnorrhiza*, *E. agallocha* and *P. paludosa*, TF was recorded as maximum from site 6 (46.42 mg g⁻¹ in *E. agallocha*, 43.55 mg g⁻¹ in *B. gymnorrhiza* and 39.54 mg g⁻¹ in *P. paludosa*) over the control plants and the increment was consistent as the salinity rise from site 1 to site 6 (Fig. 2B). But in other two taxa, these trends were not found with the raised soil salinity. In *H. fomes*, the increment of TF recorded over the control environment was up to site 3, beyond this it gradually decreased (highest in site 3, 26.72 mg g⁻¹ in site 4, 5 and 6 (Fig. 2B). The highest TF occurrence in *X. granatum* was recorded from sites 3 and 4 over control plants (almost same in the two sites, 16.54 and 16.59 mg g mg g⁻¹). It gradually came down with the soil salinity increase in site 5 and 6 (14.39 and 9.57 mg g mg g⁻¹ respectively).

To determine the free radical scavenging as an index of antioxidant ability of the investigated taxa, DPPH assay was performed. In this experiment, the percent inhibitions were increased as the substrate salinity increased in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* over the mesophytic individuals. But in *H. fomes* and *X. granatum*, the percent inhibition decreased beyond the site 4 (Fig. 2C). IC₅₀ values were calculated in all the cases which reflected the usual reverse diagrams (Fig. 2D). The highest percent inhibition was recorded in *E. agallocha* (17.8) grown in highest substrate salinity level. The magnitude of inhibition in *H. fomes* and *X. granatum* observed in site 4 (6.248 and 4.865 respectively), where in other three taxa in the same salinity level were 15.23 (*B. gymnorrhiza*), 9.87 (*E. agallocha*) and 8.64 (*P. paludosa*) and beyond this, the percent inhibition increased further for these three taxa as the salinity increased.

Antioxidant ability was further studied in all taxa with ABTS+ radical scavenging efficiency along the substrate salinity gradients. In this assay, similar results were obtained as DPPH assay in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa*, the percent inhibition of cation radical were enhanced as the salinity level hiked. But this trend was discontinues at site 4 in *H. fomes* and at site 5 in *X. granatum*, where beyond these sites along the salinity

gradient, the percent inhibition dwindled. Highest inhibition occurred at site 6 as 34.29 (*E. agallocha*), 31.95 (*P. paludosa*) and 25.26 (*B. gymnorrhiza*). In the same site *H. fomes* and *X. granatum* showed 18.84 and 23.26 percent of inhibition (Fig. 2E). Incidence of 50% inhibition (IC_{50}) depicted reverse of percent scavenging (Fig. 2F).

The Fe_2^+ chelating assay determines the antioxidants confined in the samples as reductants in colorimetric reaction, which is redox-linked and the results reveal the antioxidants reducing power. A higher absorbance tallies to a higher ferric reducing ability and IC_{50} values were calculated, which denotes the concentration of samples required for scavenging 50% of the free radicals. The present experiment publicized that in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* the reducing ability were distinct along with the increasing salinity, though in *P. paludosa* the increment was not as high as the other two. The percent inhibitions were noted from these three taxa maximized in site 6 (35.86 in *B. gymnorrhiza*, 36.85 in *E. agallocha* and 21.98 in *P. paludosa*). But *H. fomes* touched peak in site 2 (18.39), after that the value reduced as salinity increased and in last two sites, remain more or less the same. *X. granatum* exhibited its highest ferric reducing ability at site 2 (17.66), it reduced at next 4 sites with increased salinities. The graphical representation of percent inhibition was depicted in Fig.2G and the bar diagram of IC_{50} represented the reverse of the percent inhibition diagram (Fig. 2H).

A two-tailed bivariate correlation coefficient (Pearson coefficient) was calculated (using SPSS 12.0 software) for each parameter in respect to salinity (Table 2). In this part, amount of increment of total phenols and flavonoids, percent inhibition of free radical scavenging by DPPH and ABTS assay and percent ability to Fe_2^+ chelation in respect to salinity hike were considered. The analysis revealed that all studied taxa showed fairly positive correlation with the salinity, the correlation values were significant ($p \leq 0.01$) in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* ($r^2=0.974$, 0.944 and 0.949 respectively), but in case of *H. fomes* and *X. granatum*, the correlation values were 0.822 and 0.860 ($p \leq 0.05$). In case of flavonoids, significant correlation only exist in former three species (0.889, $p \leq 0.05$ in *B. gymnorrhiza*; 0.952, $p \leq 0.01$ in *E. agallocha* and 0.904, $p \leq 0.05$ in *P. paludosa*), other two have no significant relationships.

Percent inhibition of free radicals by DPPH and ABTS also exhibited the significant relationship with salinity in case of former three species. These three species showed high significant correlation in ferric ion chelation ability with salinity, whereas rest two did not reveal significant correlation

values. Percent inhibition of DPPH radical is well correlated with the incidence of TP and TF in all cases at significant level $p \leq 0.01$, except *H. fomes* ($p \leq 0.05$). In *X. granatum*, DPPH and flavonoids occurrence was not significantly correlated. ABTS assay of free radical scavenging and occurrence of TP and TF also produced the same trend for *X. granatum*. Ferric ion chelating ability in respect to TP revealed maximum correlation in *B. gymnorrhiza*, *E. agallocha*, *X. granatum* and *P. paludosa* ($p \leq 0.01$), but *H. fomes* showed least. In case of TF, except *H. fomes* and *X. granatum*, all other studied taxa were significantly correlated.

Table 2. Correlations among salinity and other investigated factors for each plant

	<i>B. gymnorrhiza</i>	<i>E. agallocha</i>	<i>H. fomes</i>	<i>P. paludosa</i>	<i>X. granatum</i>	Treatments
Salinity	0.974**	0.944**	0.822*	0.949**	0.860*	Phenols
	0.889*	0.952**	0.329	0.904*	0.294	Flavonoids
	0.928**	0.856*	0.582	0.981**	0.414	ABTS
	0.946**	0.890*	0.646	0.884*	0.734	DPPH
	0.966**	0.962**	0.316	0.919**	0.738	Fe ²⁺
DPPH	0.831*	0.968**	0.892*	0.922**	0.956**	Phenols
	0.944**	0.934**	0.864*	0.963**	0.596	Flavonoids
ABTS	0.993**	0.978**	0.868*	0.981**	0.870*	Phenols
	0.971**	0.942**	0.865*	0.959**	0.544	Flavonoids
Fe ²⁺	0.989**	0.963**	0.733	0.921**	0.975**	Phenols
	0.963**	0.986**	0.736	0.852*	0.575	Flavonoids

** Correlation is significant at the 0.01 level (2-tailed).

* Correlation is significant at the 0.05 level (2-tailed)..

The present work demonstrated the amount of total phenols, flavonoids and free radical scavenging ability of five mangroves, among which three are abundantly growing in Sundarbans region and while the other two exhibited scarce occurrence from five different islands of varying substrate salinity. The same plants were taken from mesophytic conditions, considered as control experiment. Incidence of total phenol in all studied taxa has a strong positive association with the salinity though this affinity was less significant in two taxa, *H. fomes* and *X. granatum*. TP increment was observed up to a certain limit of salinity, beyond which it decreased. In mangrove species, normally total phenols are copious components, which avert damage from herbivores (Feller *et al.*, 1999; Hernes *et al.*, 2001), but basically their major role in plant system is involve in antioxidative radical scavenging (Banerjee *et al.*, 2008; Rahim *et al.*, 2008; Zhang *et al.*, 2010).

Phenolic compounds are commonly found in plants and have been reported towards several biological activities including a strong antioxidant activity (Duh *et al.*, 1999; Chandini *et al.*, 2008). The





Fig.2. A – H. Graphical representation of different parameters with error bars of the investigated taxa collected from different sites. A. Total Phenols; B. Total Flavonoids; C - D. Percent inhibition of DPPH radical (C), IC₅₀ values of the taxa at different sites (D); E – F. Percent inhibition of ABTS radical (E), IC₅₀ values of the at different sites (F); G – H. Ferrous ion chelating ability by the extractants (G), IC₅₀ values of the taxa at different sites.

Continuation of Figure 2



antioxidant capacity of phenolic extracts is very often attributed to their radical scavenging ability mediated by hydroxyl groups (Hatano *et al.*, 1989).

The flavonoids have been recognized as important components in Plant leaves. Metcalfe (1987) opined that 20% of the carbon that have been fixed by photosynthesis have channelized through phenylpropanoid pathway leading to generate most of the naturally occurring polyphenols. It is established that plant polyphenols and, especially, flavonoids are the most important defensive antioxidants in stressful condition (Agati and Tattini, 2010; Pollastri and Tattini, 2011; Brunetti *et al.*, 2013). Agati *et al.* (2012) commented that flavonoids effectively absorb the active solar wavelengths (UV-B and UV-A), hinder the generation of reactive oxygen species (ROS) and scavenge ROS once they are formed. With several endogenous antioxidants, flavonoids play a role in optimum protection from oxidative stress caused by the increased level of ROS. Antioxidative mechanism of flavonoids includes suppressing ROS formation (by inhibition of enzymes or chelating trace elements involving ROS generation), direct ROS scavenging and up-regulating or protecting antioxidant defenses (Middleton and Kandaswami, 1993; Pietta, 2000). In the present observation, incidence of flavonoids in the leaves extracts from the three species (*B. gymnorhiza*, *E. agallocha* and *P. paludosa*) were high than those of the two (*H. fomes* and *X. granatum*). As the substrate salinity elevated in different sites, the flavonoids occurrence also increased in the former three species indicating their better suitability in increased salinity, whereas, the latter two species show different trend. The flavonoids increment over the control plants was continued up to a certain limit of substrate salinity in these two species, beyond that, its diminution designating their weaker adaptability in high saline environment. Scientific suggestion revealed that, under abiotic stress, excess generation of ROS, such as superoxide, hydroxyl and peroxy radicals occurs and the balance between antioxidation and oxidation is supposed to be a critical perception for keeping a well biological system (Davis, 2000).

It has been demonstrated that salt stress is expected to encourage the generation of ROS in plants (M'Rahs *et al.*, 2007). Plants respond to salt stress by decreasing stomatal conductance to avoid excessive water loss. This in turn decreases the internal CO₂ concentrations and slows down the reduction of CO₂ by Calvin cycle. This response leads to depletion of the oxidized NADP⁺ which acts as a final acceptor of electrons in PSI, and alternatively increases the leakage of electrons to O₂ forming O⁻² (Hsu and

Kao, 2003). Furthermore, Na^+/Cl^- toxicity resulting from salt stress could disrupt the photosynthetic electron transport and provoke electron leakage to O_2 . Nychas *et al.* (2003) claimed that polyphenols are antioxidants (having redox properties), potentially act as reducing agents, hydrogen donors and singlet oxygen quenchers. Jithesh *et al.* (2006) commented that in increased salinity, physiological water deficit causing abscisic acid regulated stomatal closure in leaves leading to limited CO_2 availability, ultimately over reduction of electron transport chain, and finally generation of ROS. In this work, free radical scavenging ability also evaluated by percent reduction of leaves extracts by DPPH, ABTS and chelation of ferric ion and were well correlated with the increasing soil salinity, at least in those species, which are predominant in *in situ* condition (*B. gymnorhiza*, *E. agallocha* and *P. paludosa*), but these trends were interrupted in *H. fomes* and *X. granatum* – the taxa which are now precarious in nature. Mangroves are inhabitants in extreme inimical microclimate with high salinity, irradiation and depleted nutrition with rapid tidal fluctuation (Naskar and Guha Bakshi, 1995). These leading to production of excess ROS and to nullify the deleterious consequences, amount of antioxidative enzymes and secondary metabolites are also sufficiently elevated. Scientific evidences proved that mangroves are generously endowed with polyphenol compounds (phenols and flavonoids), which help to cope with the oxidative stress by exploitation of potent antioxidant (Banerjee *et al.*, 2008; Chandini *et al.*, 2008).

Beside the genetic regulator, a biochemical mechanism, which are regulated in a way that allows halophytes to nurture and flourish in salt stress environment contributing them a definite advantage over other glycophytes. Inductions of antioxidants (enzymes and secondary metabolites) were shown to protect halophytes against ROS, leading to prevention of lipid peroxidation during salt stress (Hasegawa *et al.*, 2000a, b; Jithesh *et al.*, 2006). This confirms that these antioxidants are indispensable towards adaptive defense mechanism against salt stress in halophytes. In the present work, it publicized that the salinity plays a major role in occurrence of TP and TF in all taxa, except the two (*H. fomes* and *X. granatum*), where up to a certain limit of substrate salinity, TP and TF amplified, further than which, it dropped down indicating their less ability to salt adaptability. The antioxidative response in plants is complex and scavenging of ROS takes place by interaction of enzymatic and non-enzymatic components. Jithesh *et al.* (2006) also pronounced that most salt tolerance studies have explored antioxidant status in leaves. This is not

surprising considering the fact that salt stress aggravates ROS mainly in the chloroplast compartments of leaf cells. It has long been established that antioxidant ability of polyphenols and flavonoids is attributed to extent of free radical scavenging (Hatano *et al.*, 1989). Present investigation confirmed that efficient free radical scavenging aptitude signify the efficacious existence of plant taxa in salt stress. The substrate salinity of Indian Sundarbans is presently much higher and imposing much inhospitality to some key species of this area for their sustainable existence (Haq *et al.*, 1999; Nandy *et al.*, 2007). Accumulation of elevated and positively correlated antioxidative enzymes and other compatible metabolite (low molecular proteins and amino acids) have demonstrated as combat force against salinity stress in *B. gymnorrhiza*, *E. agallocha* and *P. paludosa* along with some other predominant flora of this area, but *H. fomes* and *X. granatum* are lagging behind in these regards (Dasgupta *et al.*, 2013). The present opinion on relative incidence of antioxidative secondary metabolites and ROS scavenging ability with respect to substrate salinity would be symptomatic towards the emaciated occurrence of *H. fomes* and *X. granatum* than those of the other three investigated taxa.

ACKNOWLEDGMENT

The authors are sincerely indebted to the Director, Sundarbans Biosphere Reserve and Chief Principal Conservator of Forest and Wildlife, Government of West Bengal, for providing required permission to conduct field work in the Sundarbans Reserved Forest time to time.

REFERENCES

- AGATI, G., M. TATTINI. 2010. Multiple functional roles of flavonoids in photoprotection. *New Phytol.* 186: 786-793.
- AGATI, G., E. AZZARELLO, S. POLLASTRI, M. TATTINI. 2012. Flavonoids as antioxidants in plants: Location and functional significance. *Plant Sci.* 196: 67-76.
- ALAM, E.A. 2011. Phytochemical screening on different plant parts of some succulent plants of Egypt. *New York Sci. J.* 4: 15-18.

- ALONGI, D.M. 2009. *The Energetics of Mangrove Forests*. Springer, Dordrecht, 216 p.
- ARNAO, M.B. 2000. Some methodological problems in the determination of antioxidant activity using chromogen radicals: a practical case. *Trends Food Sci. Technol.* 11: 419-421.
- BANDARNAYAKE, W.M. 2002. Bioactivities, bioactive compounds and chemical constituents of mangrove plants. *Wetl. Eco. Manag.* 10: 421-452.
- BANERJEE, D., S. CHAKRABORTY, A.K. HAZRA, S. BANERJEE, J. ROY, B. MUKHERJEE. 2008. Antioxidant activity and total phenolics of some mangroves in Sundarbans. *Afr. J. Biotech.* 7: 805-810.
- BANERJEE, L.K. 1999. *Mangroves of Orissa coast and their ecology*. Bishen Singh Mohendra Pal Singh, Dehra Dun, India. pp 41.
- BRUNETTI, C., M. DI FERDINANDO, A. FINI, S. POLLASTRI, M. TATTINI. 2013. Flavonoids as Antioxidants and Developmental Regulators: Relative Significance in Plants and Humans. *Internatl. J. Mol. Sci.* 14: 3540-3555.
- CHANDINI, S.K., P. GANESAN, N. BHASKAR. 2008. *In vitro* antioxidant activities of three selected brown seaweeds of India. *Food Chem.* 107: 707-713.
- COTELLE, N., J. L. BERNIER, J. P. CATTEAU, J. POMMERY, J. C. WALLET, E. M. GAYDOU. 1996. Antioxidant properties of hydroxy-flavones. *Free Rad. Biol. Med.* 20: 35-43.
- CUSHNIE, T.P.T., A. J. LAMB. 2005. Antimicrobial activity of flavonoids. *Internatl. J. Antimicrob. Agents.* 26: 343-356.
- DASGUPTA, N., P. NANDY, S. DAS. 2013. Salt Stress: A biochemical and physiological adaptation of some Indian halophytes of Sundarbans. In: Rout, G.R., A. B. Das (eds.) *Molecular Stress Physiology of Plants*, 155-177p. Springer, India.
- DASGUPTA, N., P. NANDY, C. SENGUPTA, S. DAS. 2012. Protein and enzymes regulations towards salt tolerance of some Indian mangroves in relation to adaptation. *Trees: Struc. Func.* 26: 377-391.

- DAVIES, K.J.A. 2000. Oxidative stress, antioxidant defenses and damage removal, repair and replacement systems. *Internat. Union of Biochem. Mol. Biol.* 50: 279-289.
- DIPLOCK, A.T. 1997. Will the 'good fairies' please prove to us that vitamin E lessens human degenerative of disease? *Free Rad. Res.* 27: 511-532.
- DUAN, X.J., W. W. ZHANG, X. M. LI, B. G. WANG. 2006. Evaluation of antioxidant property of extract and fractions obtained from a red alga, *Polysiphonia urceolata*. *Food Chem.* 95: 27-43.
- DUH, P.D., Y.Y. TU, G. C. YEN. 1999. Antioxidant activity of water extract of Harnng Jyur (*Chrysanthemum morifolium* Ramat). *Lebnesmittel-wissenschaft and Technologie* 32: 269-277.
- FOOD AND AGRICULTURAL ORGANIZATION, United Nations (FAO). 2007. *The World's Mangroves 1980 – 2005*. FAO Forestry Paper 153, FAO, Rome.
- FELLER, I.C, D. F. WHIGHAM, J. P. O'NEILL, K. L. MCKEE. 1999. Effects of nutrient enrichment on within-stand cycling in a mangrove forest. *Ecol.* 80: 2193–2205.
- GIRI, C., E. OCHIENG, L. L. TIESZEN, Z. ZHU, A. SINGH, T. LOVELAND, J. MASEK, N. C. DUKE. 2011. Status and distribution of mangrove forests of the world using earth observation satellite data. *Global Ecology. Biogeogr.* 20: 154-159.
- HALLIWELL, B., C. E. CROSS. 1994. Oxygen-derived species: their reaction to human disease and Environmental stress. *Environ. Health Persp.* 102(10): 5-12.
- HAQ, S., Z. KARIM, M. ASADUZZAMAN, F. MAHTAB. 1999. *Vulnerability and Adaptation to Climate Change for Bangladesh*. Kluwer Academic Publishers, The Netherlands. ISBN-7923-5536-9.
- HARO-VICENTE, J.F., C. MARTÍNEZ-GRACIÁ, G. ROS. 2006. Optimisation of in vitro measurement of available iron from different fortificants in citric fruit juices. *Food Chem.* 98: 639-648.
- HASEGAWA, P. M., R. BRESSAN, J. M. PARDO. 2000a. The dawn of plant salt tolerant genetics. *Trends Plant Sci.* 5: 317–319.

- HASEGAWA, P.M., R. A. BRESSAN, J. K. ZHU, H. J. BOHNERT. 2000b. Plant cellular and molecular responses to high salinity. *Ann. Rev. Plant Physiol. Plant Mol. Biol.* 51: 463-499.
- HATANO, T., R. EDAMATSU, M. HIRAMATSU, A. MORI, Y. FUJITA, T. YASUHARA, T. YOSHIDA, T. OKUDA. 1989. Effect of the interaction of tannins with co-existing substances. VI. Effect of tannins and related polyphenols on superoxide anion radical and on 1,1-diphenyl-2-picrylhydrazyl radical. *Chem. Pharma. Bull.* 37: 2016-2021.
- HERNES, P. J., R. BENNER, G. L. COWIE, M. A. GONI, B. A. BERGAMASCHI, J. I. HEDGES. 2001. Tannin diagenesis in mangrove leaves from a tropical estuary: A novel molecular approach. *Geochimica et Cosmochimica Acta.* 65: 3109-3122.
- HSU, S.Y., C. H. KAO. 2003. Differential effect of sorbitol and polyethylene glycol on antioxidant enzymes in rice leaves. *Plant Growth Regul.* 39: 83-90.
- HUDA-FAUJAN, N., A. NORIHAM, A. S. NORRAKIAH, A. S. BABJI. 2009. Antioxidant activity of plants methanolic extracts containing phenolic compounds. *Afr. J. Biotechnol.* 8: 484-489.
- IMLAY, J.A. 2003. Pathways of oxidative damage. *Ann. Rev. Microb.* 57: 395-408.
- JIA, Z.S., M. C. TANG, J. M. WU. 1999. The determination of flavonoid contents in mulberry and their scavenging effects on superoxide radicals. *Food Chem.* 64: 555-559.
- JITHESH, M.N., S. R. PRASHANTH, K. R. SIVAPRAKASH, A. K. PARIDA. 2006. Antioxidative response mechanisms in halophytes: their role in stress defence. *J. Genet.* 85: 237-254.
- KRISHNAMOORTHY, M., J. M. SASIKUMAR, R. SHAMNA, C. PANDIARAJAN, P. SOFIA, B. NAGARAJAN. 2011. Antioxidant activities of bark extract from mangroves, *Bruguiera cylindrica* (L.) Blume and *Ceriops decandra* Perr. *Indian J. Pharma.* 43: 557-562.
- LEE, HUI-WEN., P. HAKIM, A. RABU, H. A. SANI. 2012. Antidiabetic effect of *Gynura procumbens* leaves extracts involve modulation of hepatic carbohydrate metabolism in streptozotocin-induced diabetic rats. *J. Medicinal Plant Res.* 6: 796-812.

- MCIVOR, A.L., T. SPENCER, I. MÖLLER, M. SPADLING. 2012. Storm Surge Reduction by Mangroves. Natural Coastal Protection Series: Report 2. Cambridge Coastal Research Unit Working Paper 41. p.36.
- MEIR, S., J. KANNER, B. AKIRI, S. PHILOSOPH-HADAS. 1995. Determination and involvement of Aqueous Reducing Compounds in Oxidative Defence Systems of Various Senescing Leaves. *J. Agric. Food Chem.* 43: 1813-1819.
- METCALFE, R.L. 1987. Plant volatiles as insect attractants. *Crit. Rev. Plant Sci.* 5: 251-301.
- MIDDLETON, E. JR., C. KANDASWAMI. 1993. *The Flavonoids*. In: Harborne, J.B. (ed.) *Advances in Research Since 1986*, 619-652 p. Chapman and Hall, London.
- M'RAHS, Z. OUERGHI, F. EYMERY, P. REY, M. HAJJI, C. GRIGNON, M. LACHAAL. 2007. Efficiency of Biochemical protection against toxic effects of accumulated salt differentiates *Thellungiella halophila* from *Arabidopsis thaliana*. *J. Plant Physiol.* 164: 375-384.
- NANDY, P., N. DASGUPTA, S. DAS. 2009. Differential expression of physiological and biochemical characters of some Indian mangroves towards salt tolerance. *Physiol. Mol. Biol. Plants.* 15: 151-160.
- NANDY, P., S. DAS, M. GHOSE, R. SPOONER HART. 2007. Effects of salinity on photosynthesis, leaf anatomy, ion accumulation and photosynthetic nitrogen use efficiency in five Indian mangroves. *Wetl. Ecol. Manag.* 15: 347-357.
- NASKAR, K.R., D. N. GUHA BAKSHI. 1995. *Vegetation pattern of the Sundarbans. Mangrove swamps of the Sundarbans – An ecological perspective*. Naya Prakash, Calcutta, India.
- NYCHAS, G. J. E., C. C. TASSOU, P. SKANDAMIS. 2003. *Antimicrobials from herbs and spices*. In: Roller, S.M. (ed.) *Natural antimicrobials for the minimum processing of foods*, 176-200 p. CRS Press, Woodhead Publishers, NY.
- OSTROWSKAA, J., W. ŁUCZAJA, I. KASACKA, A. RÓŻAŃSKIC, S. E. ELŻBIETA. 2004. Green tea protects against ethanol-induced lipid peroxidation in rat organs. *Alcohol.* 32: 25-32.

- PARANI, M., M. LAKSHMI, B. ZEIGENHAGEN, M. FLADUNG, P. SENTHILKUMAR, A. PARIDA. 2000. Molecular phylogeny of mangroves VII. PCR-RFLP of trnS – pbsC and rbcL gene regions in 24 mangrove and mangrove associate species. *Theor. Appl. Genet.* 100: 454-460.
- PIETTA, P. G. 2000. Flavonoids as antioxidants. *J. Nat. Prod.* 63: 1035-1042.
- POLLASTRI, S., M. TATTINI. 2011. Flavonols: Old compound for old roles. *Annal. Bot.* 108: 1225–1233.
- RAHIM, A. A., E. ROCCA, J. STEINMETZ, K. M. JAIN, M. SANI IBRAHIM, H. OSMAN. 2008. Antioxidant activities of mangrove *Rhizophora apiculata* bark extracts. *Food Chem.* 107: 200-207.
- RAVIKUMAR, S., M. GNANADESIGAN, P. SUGANTHI, A. RAMALAKSHMI. 2010. Antibacterial potential of chosen mangrove plants against isolated urinary tract infectious bacterial pathogens. *Int. J. Med. Med. Sci.* 2: 94-99.
- RAVIKUMAR, S., G. RAMANATHAN, S. J. INBANESON, A. RAMU. 2011. Antiplasmodial activity of two marine polyherbal preparations from *Chaetomorpha antennina* and *Aegiceras corniculatum* against *Plasmodium falciparum*. *Parasitol. Res.* 108: 107-113.
- RE, R. N., A. PELLEGRINI, A. PROTEGGENTE, M. PANNALA, C. YANG, R. EVANS. 1999. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Rad. Biol. Med.* 26: 1231-1237.
- RIBEIRO, A.B., D. H. S. SILVA, V DA SILVA BOLZANI. 2002. *Antioxidant flavonol glycosides from Nectandra grandiflora* (Lauraceae). *Eclet. Quim. Versa Oimpressa*: 27 (special) Sao Paulo. (ISSN 0100-4670).
- SHAHIDI, F. 2000. Antioxidants in food and food antioxidants. *Food / Nahrung* 44: 158-163.
- SINGLETON, V. L., A. ROSSI. 1965. Colorimetry of Total Phenolics with Phosphomolybdic Phosphotungstic Acid Reagents. *Am. J. Enol. Viticult.* 16: 144-158.
- TIWARI, A. K. 2001. Imbalance in antioxidant defense and human diseases: Multiple approach of natural antioxidant therapy. *Curr. Sci.* 8: 1179-1187.

- UPADHYAY, V. P., RAJIV RANJAN, J. S. SINGH. 2002. The Human Mangrove Conflicts - The way out. *Curr. Sci.* 83: 1328-1336.
- VADLAPUDI, V., K. C. NAIDU. 2009. Evaluation of Antioxidant potential of selected Mangrove Plants. *J. Pharma. Res.* 2: 1742-1745.
- YILDIRIM, A., A. MAVI, A. KARA. 2001. Determination of antioxidant and antimicrobial activities of *Rumex crispus* L. extracts. *J. Agric. Food Chem.* 49: 4083–4089.
- YONGXIN, LI., YU. SHANJIANG, L. DONG, P. PETER, W. LIN. 2012. Inhibitory effects of polyphenols toward HCV from the mangrove plant *Excoecaria agallocha* L. *Bioorg. Med. Chem. Let.* 22:1099-1102.
- ZHANG, L.L., Y. M. LIN, H. C. ZHOU, S. D. WEI, J. H. CHEN. 2010. Condensed tannins from mangrove species *Kandelia candel* and *Rhizophora mangle* and their antioxidant activity. *Molecules.* 15: 420–431.
- ZURINA, H., M. F. YAM, A. MARIAM, P. AHMAD, M. Y. PAUZI. 2010. Antidiabetic properties and mechanism of action of *Gynura procumbens* water extract in streptozotocin-induced diabetic rats. *Molecules*, 15: 9008-9023.